were interrogated to identify pathogens. A host transcriptomic MLC was developed from human RNA transcripts using 70 cases. The MLC and mNGS reporting thresholds were then tested on 108 blinded cases within the cohort.

Results mNGS was 75% concordant (27/36) for detecting TB in definite TBM cases and 59% concordant (30/51) in definite/probable TBM combined. 3 TB and 3 non-TB pathogens were detected in the probable TBM group. In the possible TBM/indeterminant groups, mNGS identified 3 cases of TBM and 17 other pathogens. The combined mNGS and host-MLC displayed 83.3% (5/6) sensitivity, 86.8% (59/68) specificity, with an area under the ROC curve of 0.83 ($p=0.009$).

Conclusion mNGS identified an array of infectious TBM mimics, including many treatable and vaccine preventable pathogens. mNGS was 75% concordant with definite TBM. We further enhanced the sensitivity of the CSF mNGS assay by developing the first CSF-based host MLC to discriminate between TBM and its mimics

039 PREVALENCE OF MRI SIGNS OF INTRACRANIAL HYPERTENSION AND THEIR ASSOCIATION WITH PAPILLEDEMA: A PROSPECTIVE STUDY USING OCULAR FUNDUS PHOTOGRAPHY

^{1,2,3}Benson S Chen, ¹Benjamin I Meyer, ¹Amit M Saindane, ¹Beau B Bruce, ¹Nancy J Newman, ¹Valérie Biousse. ¹Emory University School of Medicine, Atlanta, GA, USA; ²Department of Clinical Neurosciences, University of Cambridge, Cambridge, UK; ³Cambridge Eye Unit, Addenbrooke's Hospital, Cambridge, UK

10.1136/bmjno-2021-ANZAN.39

Objectives MRI signs of intracranial hypertension (MRI-IH) are classically associated with idiopathic intracranial hypertension (IIH), but also detected in asymptomatic individuals without papilledema. The objective of this study was to determine the prevalence of MRI-IH in consecutive outpatients undergoing brain MRI for any clinical indication, and explore their association with papilledema.

Methods Prospective cross-sectional observational study of outpatients undergoing brain MRI, with ocular fundus photographs taken concurrently. Radiologic studies were analyzed for MRI-IH. Univariate analysis with Fisher's exact test or t-test was performed.

Results Of 296 patients included, the most common indication for MRI was surveillance of a brain neoplasm (27.7%). Investigation of headaches (8.8%) or disorders of raised ICP (1.4%) were uncommon. At least one MRI-IH was present in 49% of patients [empty sella (33.1%), enlarged Meckel's cave (15.9%), increased peri-optic CSF (10.8%), optic nerve tortuosity (7.8%), scleral flattening (0.7%), cephaloceles (1.4%)]. Bilateral transverse venous sinus stenosis (TVSS) was present in 3.0% of 198 patients. Five patients (1.7%) had papilledema. Compared to patients without papilledema, those with papilledema had significantly higher BMI and prior history of IIH, and increased prevalence of empty sella, optic nerve tortuosity, and TVSS on MRI. The prevalence of papilledema increased from 2.8% among patients with at least one MRI-IH to 40% among patients with four or more MRI-IH.

Conclusion MRI-IH are common in patients undergoing brain MRI, but rarely associated with papilledema. The management of patients with incidentally detected MRI-IH likely does not require systematic lumbar puncture unless concerning symptoms or papilledema are present.

040 THE QUEST TO REDUCE STROKE TREATMENT DELAYS AT A MELBOURNE METROPOLITAN PRIMARY STROKE CENTRE OVER THE LAST TWO DECADES

^{1,2}Peter SW Park, ¹Tanya Frost, ¹Peter SY Tan, ¹Joseph Wong, ²Alun Pope, ^{1,2}Helen M Dewey, ^{1,2}Philip MC Choi. ¹Department of Neurosciences, Box Hill Hospital, Eastern Health, Box Hill, VIC, Australia; ²Eastern Health Clinical School, Faculty of Medicine, Nursing and Health Sciences, Box Hill, VIC, Australia

10.1136/bmjno-2021-ANZAN.40

Objectives To examine door-to-needle time (DNT) trends and the impact of continuous quality improvement initiatives since thrombolysis became standard of care in Australian metropolitan setting for eligible acute ischaemic stroke patients.

Methods Single-centre retrospective cohort study of consecutive patients treated with thrombolysis at high volume primary stroke centre from January 2003 to December 2019. Primary outcomes are DNT, and annual number of 'Code Stroke' activations and stroke admissions. Regression modelling for potential factors associated with DNT.

Results 1,250 patients were treated with thrombolysis over 17 years; 54% were male with a median age of 76 (interquartile ranges [IQR], 66–83). Median DNT fluctuated between 70 to 93 minutes (IQR, 55–95 to 82–120) from 2003 to 2012, reaching 60 minutes in 2013 and nadir of 47 minutes in 2014. Median DNT then decreased from 58 minutes in 2015 to 51 minutes in 2019 with progressive tightening of IQR (46–78 to 40–62). Number of patients treated within 60 minutes of hospital arrival was less than 30% between 2003–2012. This rose to an average of 63% during 2015–2018 and 71% in 2019. From 2015 to 2019, per annum number of 'Code Stroke' activations increased from 940 to 1300 while stroke admissions plateaued at 750. 'Direct-to-CT' protocol and acute stroke presence were two modifiable workflow factors independently associated with faster DNT ($P<0.001$).

Conclusion Targeted quality improvement initiatives are key to reducing treatment delays in the Australian metropolitan setting. Relative stagnation in DNT improvement is concerning and needs further investigation.

041 COGNITIVE IMPAIRMENT IN LATE ONSET EPILEPSY

Xin Zhang, Armin Nikpour, Rebekah Ahmed. Royal Prince Alfred Hospital, Camperdown, NSW, Australia

10.1136/bmjno-2021-ANZAN.41

Objective Late onset, unprovoked epilepsy patients with cognitive impairment can have complex pathophysiology.¹ Our objective was to study the characteristics and contributors of cognitive impairment in this group; and how patients with dementia could be differentiated from late onset epilepsy patients.

Methods Twenty-six patients with epilepsy, onset after 50 years of age, with new cognitive complaints and 26 patients with clinically diagnosed Alzheimer's Disease (AD) were recruited. These participants had comprehensive neuropsychological and neuroimaging assessments. A subset of 17 participants from the Epilepsy group underwent longitudinal neuropsychological assessment.

Results In the Epilepsy group, the neuropsychological profile of cognitive impairment was consistent with the foci and severity of seizure activity in 46% of participants; subcortical