Age <20 years was associated with NMDA-R-IgG and MOG-IgG1 (OR=8.11 and 7.73 respectively, $p<0.001$). Age >65 years was associated with GABAB-R-IgG, LGI1-IgG, CASPR2-IgG and ANNA1-IgG (OR=7.33, 14.98, 3.67, 14.53, $p<0.001$). Women accounted for 60% of NMDA-R-IgG (CSF) and 78% of GAD65-IgG (CSF/serum) cohorts (OR=1.32, $p=0.002$, OR=2.78, $p<0.001$, respectively). Men accounted for 62% of the LGI1-IgG cohort (OR=1.87, $p<0.001$). Age and sex interacted for NMDA-R-IgG, particularly in females <20 years (OR=7.72, $p<0.001$).

Conclusion The most frequently detected were NMDA-R-IgG, LGI1-IgG, GAD65-IgG and MOG-IgG1. Age and sex associations may suggest paraneoplastic, endocrinological or aging influences on neurological autoimmunity.

004

PREGNANCY-RELATED RELAPSE IN NATALIZUMAB, FINGOLIMOD AND DIMETHYL FUMARATE-TREATED WOMEN WITH MULTIPLE SCLEROSIS

^{1,2}Wei Z Yeh, ¹Putu A Widyastuti, ^{1,2}Anneke Van der Walt, ¹Jim Stankovich, ³Eva K Havrdova, ³Dana Horakova, ³Karolina Vodehnalova, ⁴Serkan Ozakbas, ⁵Sara Eichau, ⁶Pierre Duquette, ^{8,7}Tomas Kalincik, ⁹Francesco Patti, ¹⁰Cavit Boz, ¹¹Murat Terzi, ¹²Bassem Yamout, ¹³Jeannette Lechner-Scott, ¹⁴Patrizia Sola, ²Olga Skibina, ¹⁵Michael Barnett, ¹⁶Marco Onofri, ¹⁷Maria J Sá, ^{18,19}Pamela McCombe, ²⁰Pierre Grammond, ²¹Radek Ampapa, ²²Francois Grand'Maison, ²³Roberto Bergamaschi, ²⁴Daniele LA Spitaleri, ²⁵Vincent Van Pesch, ²⁶Elisabetta Cartechini, ²⁷Suzanne Hodgkinson, ²⁸Aysun Soysal, ²⁹Albert Saiz, ^{1,2}Melissa Gresle, ³Tomas Uher, ³⁰Davide Maimone, ³¹Recai Turkoglu, ³²Raymond MM Hupperts, ^{33,34}Maria Pia Amato, ³⁵Franco Granello, ³⁶Celia Oreja-Guevara, ³⁷Ayse Altintas, ³⁸Richard Macdonell, ³⁹Tamara Castillo-Trivino, ^{1,2}Helmut Butzkueven, ⁴⁰Raed Alroughani, ^{1,2}Vilija G Jokubaitis, ⁴¹MSBase Registry.

¹Department of Neuroscience, Central Clinical School, Monash University, Melbourne, VIC, Australia; ²Multiple Sclerosis and Neuroimmunology Unit, Department of Neurology, Alfred Health, Melbourne, VIC, Australia; ³Department of Neurology and Center of Clinical Neuroscience, First Faculty of Medicine, Charles University and General University Hospital, Prague, Czech Republic; ⁴Dokuz Eylul University, Turkey; ⁵Hospital Universitario Virgen Macarena, Spain; ⁶CHUM – Hôpital Notre Dame, Canada; ⁷Melbourne MS Centre, Royal Melbourne Hospital, Melbourne, VIC, Australia; ⁸CORE, Department of Medicine, University of Melbourne, Melbourne, VIC, Australia; ⁹Department of Medical and Surgical Sciences and Advanced Technologies, GF Ingrassia, University of Catania – AOU Policlinico-San Marco, University of Catania, Italy; ¹⁰KTU Medical Faculty Farabi Hospital, Turkey; ¹¹Mayis University, Medical Faculty, Turkey; ¹²American University of Beirut, Faculty of Medicine, Nehme and Therese Multiple Sclerosis Center, Beirut, Lebanon; ¹³John Hunter Hospital, New Lambton Heights, NSW, Australia; ¹⁴Neurology Unit, Azienda Ospedaliero-Universitaria of Modena, Modena, Italy; ¹⁵Brain and Mind Centre, University of Sydney, Sydney, NSW, Australia; ¹⁶Univ G.d'Annunzio Chieti-Pescara, Italy; ¹⁷Department of Neurology, São João University Hospital Center, Porto, Portugal; ¹⁸Royal Brisbane and Women's Hospital, Herston, QLD, Australia; ¹⁹St Andrews Place, Spring Hill, QLD, Australia; ²⁰Centre de réadaptation déficience physique Chaudière-Appalache, Canada; ²¹Nemocnice Jihlava, Czech Republic; ²²Neuro Rive-Sud, Canada; ²³IRCCS Mondino Foundation, Pavia, Italy; ²⁴AORN San Giuseppe Moscati Avellino, Italy; ²⁵Cliniques Universitaires Saint-Luc, Université Catholique de Louvain, Belgium; ²⁶Ospedale Generale Provinciale Macerata, Italy; ²⁷Liverpool Hospital, Liverpool, NSW, Australia; ²⁸Bakirkoy Education and Research Hospital for Psychiatric and Neurological Diseases, Turkey; ²⁹Service of Neurology, Hospital Clinic, Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), and Institut de Neurociències, Universitat de Barcelona, Barcelona, Spain; ³⁰Centro Sclerosi Multipla, UOC Neurologia, ARNAS Garibaldi, Catania, Italy; ³¹Haydarpasa Numune Training and Research Hospital, Turkey; ³²Maaslandziekenhuis, Netherlands; ³³Department NEUROFARBA, University of Florence, Italy; ³⁴IRCCS Fondazione Don Carlo Gnocchi, Florence, Italy; ³⁵University of Parma, Italy; ³⁶Department of Neurology, Hospital Clínico San Carlos, Departamento de Medicina, Facultad de Medicina, Universidad Complutense de Madrid (UCM) and IdISSC, Madrid, Spain; ³⁷Department of Neurology, Koc University School of Medicine, Turkey; ³⁸Department of Neurology, Austin Health, Heidelberg, VIC, Australia; ³⁹Department of Neurology, Hospital Universitario Donostia, San Sebastian, Spain; ⁴⁰Amiri Hospital, Kuwait; ⁴¹MSBase Neuro-immunology Registry, Melbourne, VIC, Australia

10.1136/bmjno-2021-ANZAN.4

Objective To investigate pregnancy-related disease activity in a contemporary multiple sclerosis (MS) cohort.

Methods Data were obtained from the MSBase Registry. Term/preterm pregnancies conceived from 2011-2019 were included (modern cohort). Annualised relapse rates (ARR) were calculated before, during and after pregnancy. Predictors of intrapartum and early postpartum (1st 3 months) relapse were determined by clustered logistic and Cox regression analyses, respectively.

Results We included 1640 pregnancies from 1452 women. Disease-modifying therapy (DMT) used in the one-year pre-conception included natalizumab (n=219), fingolimod (n=147), dimethyl fumarate (DMF; n=57) and low-efficacy therapies (n=845). Preconception ARR by DMT class used before conception were: natalizumab, 0.29 (95% CI 0.22-0.37); fingolimod, 0.37 (0.28-0.49); DMF, 0.24 (0.13-0.41); low-efficacy, 0.29 (0.25-0.33); and none, 0.24 (0.19-0.31). Among women who used fingolimod or natalizumab, ARR increased during pregnancy. Intrapartum ARR decreased in preconception DMF, low-efficacy or no DMT groups. ARR spiked after delivery across all DMT groups. Natalizumab continuation into pregnancy reduced the odds of relapse during pregnancy (OR 0.76 per month [0.60-0.95], $p=0.017$). DMT re-initiation with natalizumab protected against postpartum relapse (HR 0.11 [0.04-0.32], $p<0.0001$). Breastfeeding women were less likely to relapse (HR 0.61 [0.41-0.91], $p=0.016$).

Conclusion Women with MS prescribed natalizumab or fingolimod preconception had higher rates of intrapartum and postpartum relapse. In women considered to be at high relapse risk, use of natalizumab before pregnancy and continued up to 32-34 weeks gestation, with early re-initiation after delivery is an effective option to minimise relapse risks. Strategies of DMT use have to be balanced against potential foetal/neonatal complications.

005

PSORIASIS IN MULTIPLE SCLEROSIS: AN AUSTRALIAN PREVALENCE STUDY

¹Varitsara Mangkornongsakul, ²Olivia A Charlton, ³Kevin Phan, ⁴Ariadna Fontes, ^{1,4}John Parratt, ^{1,4}Geoff Herkes, ^{5,6,7}Saxon D Smith. ¹The University of Sydney, Sydney, NSW, Australia; ²Department of Dermatology, Royal North Shore Hospital, St Leonards, NSW, Australia; ³Department of Dermatology, Liverpool Hospital, Liverpool, NSW, Australia; ⁴Department of Neurology, Royal North Shore Hospital, St Leonards, NSW, Australia; ⁵The Dermatology and Skin Cancer Centre, Gosford, NSW, Australia; ⁶Sydney Adventist Hospital Clinical School, Sydney Medical School, The University of Sydney, Sydney, NSW, Australia; ⁷Department of Dermatology, Sydney Adventist Hospital, Wahroonga, NSW, Australia

10.1136/bmjno-2021-ANZAN.5

Background Multiple Sclerosis (MS) is an immune-mediated, demyelinating disease of the central nervous system.¹ Although severe psoriasis and psoriasiform dermatitis have been noted in MS patients, the prevalence of psoriasis in these populations is uncertain and has not been explored in the Australian population.

Objectives A pilot study to estimate the prevalence of psoriasis in MS cohorts in the Australian population.

Methods A survey was conducted on 82 MS patients aged 18 and above who attended MS clinics in 2018.

Results Data was recorded for 82 patients. The mean age was 48 years for the entire cohort and 48.0 years (SD ± 11.30)