

2772

### NEW DIAGNOSIS OF MITOCHONDRIAL ENCEPHALOPATHY, LACTIC ACIDOSIS, AND STROKE-LIKE EPISODES (MELAS) PRESENTING AS BILATERAL CEREBRAL INFARCTS

<sup>1</sup>Oshi Swarup\*, <sup>1,2</sup>Nawaf Yassi, <sup>3,4</sup>Timothy Fazio, <sup>3</sup>Sarah Donoghue, <sup>1</sup>Stephen Davis. <sup>1</sup>Department of Neurology, Melbourne Brain Centre, Royal Melbourne Hospital and University of Melbourne, Melbourne, VIC, Australia; <sup>2</sup>Population Health and Immunity Division, The Walter and Eliza Hall Institute of Medical Research, Melbourne, VIC, Australia; <sup>3</sup>Metabolic Diseases Unit, Royal Melbourne Hospital, Melbourne, VIC, Australia; <sup>4</sup>Melbourne Medical School, University of Melbourne, Melbourne, VIC, Australia

10.1136/bmjno-2023-ANZAN.138

**Background** Mitochondrial disorders can manifest in stroke-like episodes which can pose several diagnostic and management challenges in the acute setting.

**Presentation** We present the case of a 45-year-old female, presenting with bilateral cerebral infarcts on a background of cerebral atrophy, cognitive decline, ataxia, sensorineural hearing loss, peripheral neuropathy, Type 1 Diabetes mellitus, and androgen insensitivity syndrome with investigations revealing a new diagnosis of Mitochondrial Encephalopathy, Lactic Acidosis, and Stroke-like episodes (MELAS).

**Management and Outcomes** Initial diagnostic investigations revealed an elevated serum and cerebrospinal fluid lactate, and MRI Brain revealed bilateral cerebral infarcts. Genetic testing on blood confirmed the presence of the m.3243A>G pathogenic variant in the mitochondrial gene *MT-TL1*, one of the common genetic mutations causing MELAS in the caucasian population. The heteroplasmy level on the sample was 24.6%. Further cascade testing identified several other affected family members. Ultimately, the patient was managed with oral arginine and subsequently oral taurine for secondary stroke prevention.

**Discussion** This case demonstrates the importance of recognising mitochondrial disorders in atypical stroke presentations as the acute and long-term treatments can vary from standard stroke care. While treatment with arginine therapy acutely is established in management of stroke-like presentations in MELAS, taurine therapy has also demonstrated potential reduction in stroke-like events in this patient group.<sup>1 2</sup> Making an accurate diagnosis is important in guiding long term management, family counselling and identifying relatives at risk who would benefit from early arginine treatment when presenting with stroke-like episodes, and screening for associated co-morbidities.

#### REFERENCES

- Rikimaru M, Ohsawa Y, Wolf AM, Nishimaki K, Ichimiya H, Kamimura N, Nishimatsu S, Ohta S, Sunada Y. Taurine ameliorates impaired the mitochondrial function and prevents stroke-like episodes in patients with MELAS. *Intern Med*. 2012;**51**(24):3351–7. doi: 10.2169/internalmedicine.51.7529. Epub 2012 Dec 15. PMID: 23257519.
- Ohsawa Y, Hagiwara H, Nishimatsu SI, Hirakawa A, Kamimura N, Ohtsubo H, Fukai Y, Murakami T, Koga Y, Goto YI, Ohta S, Sunada Y; KN01 Study Group. Taurine supplementation for prevention of stroke-like episodes in MELAS: a multi-centre, open-label, 52-week phase III trial. *J Neurol Neurosurg Psychiatry*. 2019 May;**90**(5):529–536. doi: 10.1136/jnnp-2018-317964. Epub 2018 Apr 17. PMID: 29666206; PMCID: PMC6581075.

2773

### NOVEL BAG3 VARIANT IDENTIFIED IN A CASE OF ADULT-ONSET MYOFIBRILLAR MYOPATHY

<sup>1,2</sup>Tram BT Nguyen\*, <sup>2,3</sup>Richard Lin, <sup>3,4</sup>Suzanna EL Temple, <sup>1</sup>Henry LT Vo, <sup>1</sup>Daniel S Green, <sup>5,6</sup>Anthony Cheong, <sup>7</sup>Roger Pamphlett, <sup>1</sup>Zeljka Calic. <sup>1</sup>Department of Neurology, Liverpool Hospital, Liverpool, NSW, Australia; <sup>2</sup>South West Sydney Clinical School, UNSW Sydney, Sydney, NSW, Australia; <sup>3</sup>Department of Clinical Genetics, Liverpool Hospital, Sydney, NSW, Australia; <sup>4</sup>School of Paediatrics, UNSW Sydney, Sydney, NSW, Australia; <sup>5</sup>Molecular Medicine Laboratory, Concord Repatriation General Hospital, Sydney, NSW, Australia; <sup>6</sup>Sydney Medical School, University of Sydney, Sydney, NSW, Australia; <sup>7</sup>Department of Neuropathology, Brain and Mind Centre, Royal Prince Alfred Hospital, Sydney, NSW, Australia

10.1136/bmjno-2023-ANZAN.139

**Objectives** Myofibrillary myopathies (MFM) are a heterogeneous group of disorders characterised by myofibrillar dissolution and accumulation of protein degradation products in myofibres. *BCL2-associated anthanogene 3* (*BAG3*)-related MFM has been reported to present with sensorimotor neuropathy and cardiomyopathy. We report a case of a 54-year-old man with a progressive sensorimotor axonal neuropathy with respiratory involvement who was found to have a novel *BAG3* variant.

**Case A** 54-year-old man presented with a six-month history of increasing dyspnoea with severe restrictive ventilatory dysfunction evident on pulmonary function testing. He has a five-year history of progressive sensorimotor disturbance in his limbs with muscle cramps and gait impairment leaving him wheelchair-bound. Examination revealed marked wasting and florid fasciculations in the upper and lower limbs, distal greater than proximal weakness, absent reflexes and a stocking-and-glove distribution sensory disturbance. Electrodiagnostic studies confirmed a severe length-dependent sensorimotor axonal neuropathy with active and chronic denervation on electromyography. Singleton whole exome sequencing identified a heterozygous missense variant of unknown significance in *BAG3* [NM\_004281.3:c.625C>G, p.(Pro209Ala)]. Other variants in *BAG3* at the same amino acid residue [p.(Pro209Leu) and p.(Pro209Gln)] have been reported to be associated with *BAG3*-related MFM. Subsequent muscle biopsy demonstrated chronic denervation and reinnervation of myofibres with pseudodystrophic changes, suggestive of an underlying myofibrillar myopathy. He was managed supportively with nocturnal bilevel positive airway pressure and discharged into supported accommodation.

**Conclusion** This is a case of adult-onset myofibrillary myopathy suspected to be due to a novel missense *BAG3* variant located in a previously reported *BAG3* mutation hotspot (Proline 209).

2774

### PLANNED DOSE REDUCTION OF OCRELIZUMAB IN RELAPSING-REMITTING MULTIPLE SCLEROSIS: A SINGLE CENTRE OBSERVATIONAL STUDY

<sup>1</sup>Trung Tran\*, <sup>2</sup>Mike Boggild. <sup>1</sup>James Cook University, Townsville, QLD, Australia; <sup>2</sup>Department of Neurology, Townsville University Hospital, Townsville, QLD, Australia

10.1136/bmjno-2023-ANZAN.140